

An analysis of rockfish landings associated with commercial salmon troll fishery off California

Don Pearson

Steve Ralston

Eric Bjorkstedt

National Marine Fisheries Service
Southwest Fisheries Science Center
Fisheries Ecology Division

Presented to the MLPA Master Plan Science Advisory
Team on January 8, 2007

Data & Analysis (1 of 2)

- Data:
 - Landing Receipts (1997-2005)
 - California Cooperative Survey (Port Sampling)
- Trip: all landings by a given boat on a given day
 - accounts for many but not all split trips (catches from same trip might be landed on different days)
- ‘Salmon trip’: a trip on which salmon were landed, regardless of whether salmon were main catch

Data & Analysis (2 of 2)


- 99% of all salmon landed by troll
 - trips with landings based on all other gear types dropped from analysis
- 99.9% of all salmon landings in ‘Chinook salmon’ market category
 - trips with landings based on all other market categories dropped from analysis
- Compiled associated landings in all rockfish market categories
- Stratified by port complex
 - Crescent City, Eureka, Fort Bragg, [Bodega Bay](#), [San Francisco](#), Monterey, Morro Bay, and Santa Barbara

Commercial Salmon Troll-Associated Rockfish Landings (CSTARL)



Port Complex	Rockfish (lbs)	Number of Salmon Trips	Bycatch Rate (lbs rockfish/trip)
Crescent City	579	516	1.12
Eureka	2,999	2,715	1.10
Fort Bragg	17,442	8,428	2.07
Bodega Bay	45,415	15,597	2.91
San Francisco	55,391	21,169	2.62
Monterey	27,041	25,742	1.05
Morro Bay	35,446	4,423	8.01
Santa Barbara	1,645	516	3.19
Combined	185,958	79,106	2.35

Mean CSTARL <3 lbs/trip in port complexes within study area over entire period 1997-2005

Most salmon trips do not land rockfish.



Most salmon trips that do land rockfish are dominated by salmon; exceptions are rare and more so in 2000s.




CSTARL: Species Composition

Yellowtail rockfish and chilipepper were largest components of landings associated with salmon troll fishery (~17% each).


Most other species represent between 0.5% to 5% of total CSTARL.

Species vary in likely susceptibility to commercial salmon trolling gear.



CSTARL v. Total RL

Salmon Troll Rockfish as a Percent of Total Rockfish



- Rockfish landings associated with commercial salmon troll fishery are very small component of total rockfish landings.

Closing thoughts

- Overall CSTARL is very small, regardless of metric (e.g., weight/trip, RF:Salmon, CSTARL:TRL)
- Trips with positive CSTARL are rare
- CSTARL dominated by two species: yellowtail and chilipepper
- Rockfish species' susceptibility to trolling gear should be considered in evaluating species-specific consequences of commercial salmon trolling
- Tightening rockfish regulations might contribute to trend of declining CSTARL from 1990s to 2000s