



Ecology of Military Islands

- San Clemente and San Nicolas islands are the most isolated and remote islands in the Southern California Bight
- Less accessible to various interest groups
 - Larger kelp bass, sheephead and lobsters
 - Abundant giant seabass
- Less likely to be affected by mainland pollution
- Least impacted by invasive species
 - E.g., Sargassum filicinum



Ecology of Military Islands

- More rocky reef habitat and kelp beds
 - Less pollution and sedimentation, and presence of large predators contribute to abundant kelp
- Significantly deeper kelp habitat
 - Due to water clarity
- Unusually large stands of surfgrass
- Largest populations of purple hydrocoral
 - Excluding Farnsworth Bank (Santa Catalina)
- Critical and substantial marine mammal haulouts and bird roosting areas



Ecological Features of San Clemente

- Northern range extensions of Panamic species
 - Panamic arrow crab (Stenorhynchus debilis)
 - Warty sea slug (*Pleurobranchus areolatus*)
 - Arbacia sea urchin (Arbacia incisa)
 - Guadalupe cardinalfish (Apogon guadalupensis)
 - Pink cardinalfish (A. pacificus)
 - Swallow damselfish (Azurina hirundo)
 - Purple brotula (Oligopus diagrammus)
- Only rock-based morph of elk kelp (*Pelagophycus porra*) at the islands
- Remnant population of endangered white abalone (*Haliotis sorenseni*)



Ecological Features of San Nicolas

- Significantly deeper eelgrass habitat
 - Due to water clarity
- Largest remaining remnant populations of endangered black abalone
- Presence of large predators enhances ecosystem function
 - Larger predatory fish (e.g. kelp bass, sheephead, giant seabass) and invertebrates (e.g., lobster)
 - Resident southern sea otter population




Ecological Features of Begg Rock

- Emergent sheer pinnacle reef 10 miles west of San Nicolas
- Unique invertebrate assemblage, including:
 - Purple hydrocoral
 - Shallow aggregations of *Metridium* and other anemones




Habitats at San Clemente Island





Habitats at San Nicolas Island





East Channel Islands Bioregion

Proposed Military Closures

MPAs proposed by the RSG on
San Clemente Island

Draft MPA Arrays and Draft External MPA Proposals

*LA = Lapis A

*OA = Opal A

*TB = Topaz B


*XB = External B


LB = Lapis B

OB = Opal B

TA = Topaz A

XA = External A





^{*}Includes pending military closures



East Channel Islands Bioregion

Proposed Military Closures

MPAs proposed by the RSG on
San Clemente Island

Draft MPA Arrays and Draft External MPA Proposals

*LA = Lapis A

*OA = Opal A

*TB = Topaz B


*XB = External B


LB = Lapis B

OB = Opal B

TA = Topaz A

XA = External A





^{*}Includes pending military closures



East Channel Islands Bioregion

Proposed Military Closures

MPAs proposed by the RSG on
San Clemente Island

Draft MPA Arrays and Draft External MPA Proposals

*LA = Lapis A

*OA = Opal A

*TB = Topaz B


*XB = External B


LB = Lapis B

OB = Opal B

TA = Topaz A

XA = External A





^{*}Includes pending military closures



East Channel Islands Bioregion

Proposed Military Closures

MPAs proposed by the RSG on
San Clemente Island

Draft MPA Arrays and Draft External MPA Proposals

*LA = Lapis A

*OA = Opal A

*TB = Topaz B


*XB = External B


LB = Lapis B

OB = Opal B

TA = Topaz A

XA = External A






^{*}Includes pending military closures



West Channel Islands Bioregion



Draft MPA Arrays and Draft External MPA Proposals

*LA = Lapis A

*OA = Opal A

*TB = Topaz B


*XB = External B


LB = Lapis B

OB = Opal B

TA = Topaz A

XA = External A






^{*}Includes pending military closures



West Channel Islands Bioregion



Draft MPA Arrays and Draft External MPA Proposals

*LA = Lapis A

*OA = Opal A

*TB = Topaz B


*XB = External B


LB = Lapis B

OB = Opal B

TA = Topaz A

XA = External A






^{*}Includes pending military closures



West Channel Islands Bioregion



Draft MPA Arrays and Draft External MPA Proposals

*LA = Lapis A

*OA = Opal A

*TB = Topaz B


*XB = External B


LB = Lapis B

OB = Opal B

TA = Topaz A

XA = External A






^{*}Includes pending military closures



West Channel Islands Bioregion



Draft MPA Arrays and Draft External MPA Proposals

*LA = Lapis A

*OA = Opal A

*TB = Topaz B


*XB = External B


LB = Lapis B

OB = Opal B

TA = Topaz A

XA = External A





^{*}Includes pending military closures



San Nicolas Island Pending Military Closure 1

- Located in West Channel Islands Bioregion
- Includes:
 - Some kelp (Macrocystis pyrifera)
 - Low relief habitats
 - Abundant purple urchins
 - One subtidal monitoring site and one biodiversity monitoring site
- Does not include:
 - Large persistent kelp beds (to the northwest)
 - Resident breeding southern sea otter (*Enhydra lutris nereis*) population (to the northwest)



San Clemente Island Pending Military Closure 1

- Located in East Channel Islands Bioregion
- Includes:
 - Castle Rock and 9-Fathom Bank
 - Submerged rocky pinnacles
 - Abundant giant kelp (*Macrocystis pyrifera*), deeper than other areas due to water clarity
 - Large populations of purple coral (Stylaster)
 - Rare rock morph of elk kelp (Pelagophycus porra)
- Does not include:
 - Moon (or Seal) Rock (to the east), which supports a marine mammal haulout and seabird colony
 - sandy beach habitat at West Cove (to the south)



San Clemente Island Pending Military Closure 2

Includes:

- Narrow rocky shore with intermittent surfgrass beds and a few patches of eelgrass
- Narrow band of kelp parallel to the shoreline
- Seafloor drops rapidly to deeper sand

Does not include:

- Extensive eelgrass beds between White Rock and Purse Seine Rock (to the south)
- Deep water corals (Stylaster), which are abundant on the west side of San Clemente and Santa Catalina islands