

**Potential Impacts of Sex- and
Size-Selective Harvesting in Crab Populations**
Prepared for the MLPA Master Plan Science Advisory Team
Revised January 3, 2008

Several studies have shown there potentially can be negative population impacts through sex- and size-selective harvesting of large males (Hines et al. 2003). These studies have primarily been conducted using blue crabs (*Callinectes sapidus*) and snow crabs (*Chionoecetes opilio*), but it is conceivable that the changes in reproductive success and female survivorship could occur in commercially-harvested crab populations in the NCCSR as well.

Removing male crabs from populations can result in skewed sex ratios and reduced male size (Carver et al. 2005, Sainte-Marie & Lovrich 1994). These results can in turn lead to decreased copulation time (Rondeau & Sainte-Marie 2001), decreased sperm production (Kendall et al. 2002), fewer successful copulations due to exhausted sperm supplies (Carver et al. 2005), decreased female survivorship (Jivoff 1997, Sainte-Marie et al. 1999), and increased female injury (Shirley & Shirley 1988, Rondeau & Sainte-Marie 2001).

Additionally, female Dungeness crabs (*Cancer magister*) are known to preferentially mate with males that are larger than they are, leading to decreased mating opportunities for large (and more fecund) females in areas with male-biased fisheries (Smith & Jaimeson 1991a). Indeed, in areas of intense fishing pressure, large females are virtually never ovigerous due to a lack of males large enough for copulation (T. Shirley, pers. comm.). Thus removing high numbers of large males from crab populations could cause significant reductions in reproductive output and have a negative impact on female survivorship.

Recent evidence suggests that the negative impacts of removing male crabs from populations will not be tempered by the movement of crabs into and out of the fished area. Studies of *Cancer magister* in Alaska, British Columbia, and northern California have shown that adult crabs move much less than previously assumed, and could have significant site fidelity and little contact with neighboring populations (Diamond & Hankin 1985, Smith & Jaimeson 1991b, Stone & O'Clair 2001, Stone & O'Clair 2002). These studies are supported by evidence of dramatic increases in the size of male crabs in historically fished areas that are incorporated into marine protected areas (Taggart et al. 2004).

References

- Carver, A.M., T.G. Wolcott, D.L. Wolcott, and A.H. Hines. 2005. Unnatural selection: effects of a male-focused size-selective fishery on reproductive potential of a blue crab population. *J. Exper. Mar. Biol. Ecol.* 319: 29-41.
- Diamond, N. and D.G. Hankin. 1985. Movements of adult female Dungeness crabs (*Cancer magister*) in northern California based on tag recoveries. *Can. J. Fish. Aq. Sci.* 42: 919-926.
- Hines, A.H., P.R. Jivoff, P.J. Bushmann, J. vanMontfrans, D.L. Wolcott, and T.G. Wolcott. 2003. Evidence for sperm limitation in female blue crabs (*Callinectes sapidus*). *Bull. Mar. Sci.* 72: 287-310.

- Jivoff, P. 1997. The relative roles of predation and sperm competition on the duration of the post-copulatory association between the sexes in the blue crab, *Callinectes sapidus*. *Behav. Ecol. Sociobiol.* 40: 175-185.
- Kendall, M.S., D.L. Wolcott, T.G. Wolcott, and A.H. Hines. 2002. Influence of male size and mating history on ejaculates in the blue crab, *Callinectes sapidus*. *Mar. Ecol. Prog. Ser.* 230: 235-240.
- Rondeau, A. and B. Sainte-Marie. 2001. Variable mate-guarding time and sperm allocation by male snow crabs (*Chionoecetes opilio*) in response to sexual competition, and their impact on the mating success of females. *Biol. Bull.* 201: 204-217.
- Sainte-Marie, B. and G.A. Lovrich. 1994. Delivery and storage of sperm at first mating of female *Chionoecetes opilio* (Brachyura: Majidae) in relation to size and morphometric maturity of male parent. *J. Crustac. Biol.* 14: 508-521.
- Sainte-Marie, B., N. Urbani, J.-M. Sevigny, F. Hazel, and U. Kuhnlein. 1999. Multiple choice criteria and the dynamics of assortative mating during the first breeding season of female snow crab *Chionoecetes opilio* (Brachyura, Majidae). *Mar. Ecol. Prog. Ser.* 181: 141-153.
- Shirley, S.M. and T.C. Shirley. 1988. Appendage injury in Dungeness crabs *Cancer magister* in southeastern Alaska, USA. *Fish. Bull.* 86: 156-160.
- Smith, B.D. and G.S. Jamieson. 1991a. Possible consequences of intensive fishing for males on the mating opportunities of Dungeness crabs. *Amer. Fish. Soc.* 120: 650-653.
- Smith, B.D. and G.S. Jamieson. 1991b. Movement, spatial distribution, and mortality of male and female Dungeness crab *Cancer magister* near Tofino, British Columbia, Canada. *Fish. Bull.* 89: 137-148.
- Stone, R.P. and C.E. O'Clair. 2001. Seasonal movements and distribution of Dungeness crabs *Cancer magister* in a glacial southeastern Alaska estuary. *Mar. Ecol. Prog. Ser.* 214: 167-176.
- Stone, R.P. and C.E. O'Clair. 2002. Behavior of female Dungeness crabs, *Cancer magister*, in a glacial southeast Alaska estuary: homing, brooding-site fidelity, seasonal movements, and habitat use. *J. Crust. Biol.* 22: 481-492.
- Taggart, S.J., T.C. Shirley, C.E. O'Clair, and J. Mondragon. 2004. Dramatic increase in the relative abundance of large male Dungeness crabs *Cancer magister* following closure of commercial fishing in Glacier Bay, Alaska. *Amer. Fish. Soc. Symp.* 42: 243-253.