

California MLPA Master Plan Science Advisory Team
Summary of SAT Marine Bird and Mammal Evaluations for the
MLPA Blue Ribbon Task Force Recommendations as
Adopted on October 26, 2010
Revised January 17, 2011

At its meeting on October 25-26, 2010, the Marine Life Protection Act (MLPA) Blue Ribbon Task Force (BRTF) adopted seven motions with recommendations related to marine protected areas (MPAs) and special closures in the MLPA North Coast Study Region.

The first recommendation forwards the Revised Round 3 North Coast Regional Stakeholder Group (NCRSG) MPA Proposal (RNCP) and the North Coast Special Closures Recommendation to the California Fish and Game Commission. While special closures can provide significant benefits to breeding seabirds and moderate levels of protection to the breeding endangered Steller Sea Lion, overall the proposed MPAs provide few additional benefits to marine mammals and sea birds in northern California. Revisions to the Round 3 NCRSG MPA Proposal included name changes to three MPAs and updated recreational uses intended to accommodate tribal uses. Given that the initially proposed MPAs (prior to revision) provided few additional benefit to marine mammals and sea birds, the revisions do not change the proposal in terms of benefits to marine birds and mammals (see attachments A and B for Round 3 evaluation results)¹. The results of the Round 3 NCRSG MPA Proposal evaluation therefore also apply to the RNCP.

The BRTF is also forwarding the North Coast Enhanced Compliance Alternative MPA Proposal (ECA) to the California Fish and Game Commission. The ECA starts with the RNCP and then makes several modifications, including dividing four SMCAs into an offshore component and a nearshore “ribbon”, reducing the proposed allowed uses in the offshore SMCAs and maintaining the proposed allowed uses in the nearshore ribbon SMCAs. The overall geographic placement of MPAs is the same in the ECA and RNCP; the SAT Bird and Mammal Work Group determined that the differences between the ECA and RNCP are not great enough to require an additional evaluation.

The SAT emphasizes that human activity in these nearshore areas (and the adjacent offshore areas) will continue to impact marine mammals and sea birds that depend on these areas for breeding, resting and feeding. Marine mammals and sea birds would benefit from enhanced protection in the future.

¹ These evaluations can also be found on the MLPA website under the BRTF's October 25-26, 2010 meeting agenda at http://www.dfg.ca.gov/mlpa/meeting_102510.asp.

ATTACHMENT A

California MLPA Master Plan Science Advisory Team Evaluation of Benefits to Marine Birds from Round 3 North Coast Proposed Marine Protected Areas and Special Closures *October 14, 2010*

Overview

Marine birds are long-lived species, often living more than 20 years (Clapp et al. 1982) that produce few offspring and provide a large amount of parental care compared to most marine species. Thus, marine bird populations can be slow to rebound from adverse human and environmental impacts. Additionally, because marine birds feed near the top of marine food webs, are highly visible, relatively inexpensive to study and respond to oceanographic variability, they are often viewed as indicators of the marine environment (see Cairns 1992).

Marine birds can be categorized into four broad categories based on habitat use: seabirds, shorebirds, waterfowl and marsh birds. Seabirds use coastal waters and at-sea habitats; many come to land only to breed. There are, however, a number of seabird species that occur in the north coast study region (NCSR) that depend on land for resting and preening throughout the year. Shorebirds consist of multiple species of sandpipers and plovers that utilize intertidal habitat along the coast and within bays and estuaries. Waterfowl consist of ducks, grebes and loons that forage and raft in nearshore waters and within bays and estuaries. Marsh birds consist of herons and egrets that typically forage along the coasts of bays and estuaries. There are 13 species of breeding seabirds, more than 25 species of shorebirds, more than 25 species of waterfowl, and 6 species of marsh birds that use the NCSR for breeding, migration, and/or overwintering.

While marine birds are not targeted by recreational or commercial fisheries, they can benefit both directly and indirectly from marine protected area (MPA) establishment. Direct benefits include reduced disturbance at breeding and roosting sites and lower probability of interaction with humans and fishing gear at foraging areas. Indirect benefits include reduced competition for important prey resources. We conducted five separate analyses on proposed MPA arrays to estimate levels of direct and indirect benefits to marine birds: 1) protection of seabird breeding colonies and hot spots, 2) protection of major seabird roosts, 3) protection of nearshore foraging areas, 4) protection of neritic foraging 'hot spots', and 5) protection of estuary and coastal habitats and shorebirds and waterfowl within those habitats. In this document, proposed MPAs for the NCSR are evaluated for their potential benefits to marine birds. Evaluations follow the methods described in *Draft Methods Used to Evaluate MPA Proposals in the MLPA North Coast Study Region*.

Protection at Seabird Breeding Colonies, Hot Spots and Roosting Sites

Some seabird species breeding in the NCSR such as guillemots, murrelets, and petrels only come to land to breed and spend the remainder of their lives at sea. Others, such as most pelicans, cormorants and gulls, come to shore on a daily basis to rest and preen. For pelicans and cormorants, trips ashore are essential for survival because their wettable plumage must be dried to avoid hypothermia (Palmer 1962). Thus, it is important that both breeding and roosting sites be protected against human disturbances. For most species, preferred breeding and roost habitats are on offshore rocks, islands, or mainland cliffs free of mammalian predators.

Most species are known to be sensitive to human disturbance to varying degrees (summarized in Carney and Sydeman 1999). Impacts of human disturbance are known to be greatest at breeding sites, where reproduction can be dramatically affected. Because most seabirds are colonial breeders (i.e., nesting in high concentrations), high proportions of populations can be affected by severe or frequent disturbances. Impacts to birds tend to be most pronounced when humans enter the immediate area. Responses vary by species and location, but for many species, intrusion results in most if not all birds fleeing from the immediate area. Birds on nests often will flee, leaving the eggs or chicks behind. During that time, nest contents are vulnerable to predators such as gulls and ravens, exposed to the elements, and susceptible to displacement. While some birds return to nests once an intruder has gone, others tend to abandon nesting efforts. For example, Brandt's Cormorants have been observed to abandon nests en masse from even single events of human intrusion to the colony (McChesney 1997). Many studies have documented reductions in breeding success and colony attendance, as well as colony abandonment, resulting from human intrusion (Carney and Sydeman 1999).

Although often not as easily identified, activities such as close approaches to colonies and roosts or loud noises can evoke responses similar to direct human intrusions. Close approaches can include humans on foot, boats, low-flying aircraft, motor vehicles, surfers, or other sources (Jaques et al. 1996, Carney and Sydeman 1999, Jaques and Strong 2002). Studies of such disturbances on seabirds and other waterbirds have shown various results that often depend on species, location, habitat and level of habituation to human activity. However, several studies have shown reductions in breeding success or population sizes as a result of such human disturbance (e.g., Wallace and Wallace 1998, Carney and Sydeman 1999, Thayer et al. 1999, Beale and Monaghan 2004, Bouton et al. 2005, Rojek et al. 2007). In some cases, reductions in breeding success from disturbance can occur in the absence of visible behavioral changes (Beale and Monaghan 2004).

Protection of Food Resources and Foraging Areas

During the breeding season, marine birds are central place foragers, continuously returning to the breeding site throughout the day to provision young. Provisioning young is energetically taxing to breeding adults and the spatial constraints of central place foraging makes them highly dependent on localized prey availability (Pichegru et al. 2009). Marine birds may benefit from MPA establishment if there is a subsequent increase in their forage base. Prey availability has been shown to affect coloniality (whether birds form large or small colonies), the timing of reproduction, clutch sizes and levels of egg abandonment, chick growth and non-predator related chick mortality (Anderson and Gress 1984, Safina and Burger 1988, Pierotti and Annetti 1990, Massey et al. 1992, Ainley et al. 1995, Monaghan 1996, Golet et al. 2000).

We have identified two general foraging strategies used by seabirds within the NCSR: 1) nearshore foraging that occurs close to the breeding colony and 2) foraging at neritic 'hot spots' that attract congregations of pelagic prey. For our purposes, we defined nearshore foraging as a strategy used by breeding seabirds that typically forage within three miles of the colony. These species are sensitive to changes in local prey availability that can have dramatic effects on breeding success, survivorship and population status (Ainley and Boekelheide 1990, Nur and Sydeman 1999, Sydeman et al. 2001). For example, the Pelagic Cormorant and Pigeon Guillemot colonies at the Southeast Farallon Islands have undergone declines in reproductive performance and population size that are consistent with a decline in the local availability of juvenile rockfish (Sydeman et al. 2001, Warzybok and Bradley 2007).

Additionally, Robinette et al. (2007) showed that both spatial and temporal variability in sanddab recruitment was reflected in the diet of Pigeon Guillemots breeding at Point Arguello, central California. Establishing MPAs adjacent to the breeding colonies of seabirds with short foraging ranges will provide protection by decreasing competition for local prey resources and reduced displacement by boats during foraging. 'Hot spot' foraging is a strategy used by both central place foragers and migrant and overwintering birds not constrained to a breeding colony. Many studies have shown that neritic foraging seabirds congregate in predictable areas (e.g., Ford et al. 2004, Yen et al. 2004) and it has even been suggested that these congregations can be used to select areas for MPA establishment (see Harris et al. 2007, Pichegru et al. 2009). Establishing MPAs in areas of high seabird concentrations will reduce direct interactions with humans similarly targeting these areas of high prey concentrations.

Protection of Shorebirds and Wintering Waterfowl and Estuary and Coastal Habitats

Protecting the intertidal habitat of estuaries and coastal beaches will likely have direct benefits for shorebirds. For waterfowl, the eelgrass beds of the coastal estuaries provide food that is crucial for several species of geese and dabbling ducks. Additionally, waterfowl have been shown to be impacted by human caused disturbances (see Peters and Otis 2006). Protection of eelgrass beds, and estuarine habitat in general, would provide direct benefits to these birds. Finally, protecting the prey base of foraging marsh birds will provide benefits through reduced competition with humans.

Of special interest is the population of Marbled Godwits in Humboldt Bay as there is evidence that the majority of godwits wintering there are from the Alaska breeding population, which is separate from the rest of the Marbled Godwit breeding population and much smaller in numbers. High Marbled Godwit feeding densities have been documented at Samoa Bridge, Eureka Slough and the Elk River Mouth, but the Round 3 NCRSG MPA proposal does not capture these areas. The mudflats between Manila and Samoa on the west shore of Arcata Bay have higher mean densities of shorebirds than the other sites in Humboldt Bay, and again this MPA proposal does not capture this area.

Methods

Evaluations follow the methods described in the *Draft Methods Used to Evaluate Marine Protected Area Proposals in the MLPA North Coast Study Region*. Proposed MPAs would provide protection only against consumptive activities. Non-consumptive activities such as kayaking and surfing can still create disturbances at seabird breeding and roosting sites. This issue can be addressed through the use of no-entry special closure areas. Special closures are considered to provide the greatest benefit to marine birds, followed by state marine reserves (SMRs) and some state marine conservation areas (SMCAs) depending on the proposed regulations (see Table 9.2 in *Draft Methods Used to Evaluate Marine Protected Area Proposals in the MLPA North Coast Study Region* for criteria to qualify SMCAs to be included in evaluations). In Round 3, the MLPA North Coast Regional Stakeholder Group (NCRSG) came to agreement on a single marine protected area (MPA) proposal. In the NCRSG MPA Proposal, some MPAs propose recreational uses that are intended to accommodate tribal activities but are open to all non-commercial users to maintain compliance with California law.

For Round 3, we evaluated the NCRSG MPA Proposal (NCP) using standard methods and we also performed a supplemental evaluation. The standard evaluation includes all recreational take proposed in each MPA including recreational take intended only to accommodate tribal uses but open to all recreational users. The NCRSG MPA Proposal - Supplemental Evaluation (SUP) does not include proposed recreational take intended only to accommodate tribal uses. The supplemental evaluation is referred to as NCRSG MPA Proposal – Supplemental Evaluation (SUP). The NCRSG also forwarded a Round 3 NCRSG Special Closures Recommendation, which is separate from the NCRSG MPA Proposal but intended to accompany it and is, therefore, included in our evaluation. The evaluation includes analyzing the potential benefits to: 1) seabird breeding areas, 2) seabird roosting areas 3) nearshore seabird foraging areas, 4) neritic foraging areas, and 5) shorebirds and waterfowl and the estuarine waterways and coastal habitats they use.

Results

Seabird Breeding Colonies and Hot Spots

The abundance and distribution of all seabird species breeding within the NCSR are shown in Table 1. Common Murres are by far the most abundant species breeding in the NCSR, accounting for 85% of the total breeding seabirds in the NCSR.

Table 2 shows the potential benefits provided by each proposed MPA and proposed special closure in Round 3. Table 3 shows the summary of benefits for the NCRSG MPA Proposal and the NCRSG MPA Proposal – Supplemental Evaluation, based on SMRs and SMCAs meeting the criteria for this analysis, and the NCRSG Special Closures Recommendation.

The Round 3 NCRSG MPA Proposal and the NCRSG Special Closures Recommendation protect most breeding seabirds and hot spots, with approximately 65% of the breeding seabirds and 5 of the 8 designated hot spots included. This is less than the round 2 Ruby 1 proposal (85% of breeding birds protected) almost entirely due the omission of breeding hotspots at Green, Flatiron, and False Cape rocks. The NCRSG Special Closures Recommendation and NCRSG MPA Proposal together protect large numbers of the seabird species that nest on the surface in large colonies and are particularly sensitive to disturbance events such as the Brandt's Cormorant (50%) and Common Murre (70%). It also includes a high proportion of Rhinoceros Auklet (95%) and Tufted Puffin (50%) colonies. The NCRSG MPA Proposal – Supplemental Evaluation showed some additional benefits to Black Oystercatchers, Pelagic Cormorants, Pigeon Guillemots and Western Gulls. Success in protection of seabird colonies was driven by special closure designations (Tables 2 and 3). Note that special closures are based on number and species of birds, but special closure sites are not equally sensitive to vessel disturbance due to their topography. False Klamath Rock, for example, has high cliff faces below the colonies and nesting birds are not generally affected by nearby vessels, whereas Flatiron and False Cape rocks have lower relief with seabirds and pinnipeds affected by close vessels.

Major Seabird Roosts

Data on California Brown Pelican roosting abundance and distribution were used in this analysis to identify major seabird roosts. California Brown Pelicans have been well studied in the NCSR and use habitats used by other roosting seabirds. All pelican roosts were placed in

one of three categories depending on the number of pelicans observed at roost sites. Roosts were placed in the 'high' category if maximum counts exceeded 500 pelicans, 'medium' if 100-500 pelicans were observed, and 'low' if never more than 100 pelicans were observed. In the north coast study region, there are many small and medium pelican roosts and few large roosts.

Table 5 shows the number of roosts captured by all proposed MPAs and special closures while Table 6 shows the summary of number of roosts captured by MPAs meeting the criteria to provide benefits to seabirds for each evaluation and special closures. Proposal 0 did not capture any important pelican roosts in the north coast study region in qualifying MPAs (based on proposed allowed uses and criteria in Table 9.2 in *Draft Methods Used to Evaluate Marine Protected Area Proposals in the MLPA North Coast Study Region*).

The number of pelican roosts included in the NCRSG MPA Proposal are low, with 7 roosts contained in the MPA proposal, 3 in qualifying MPAs for the NCP and an additional one for the SUP, and 5 more in special closures, for a total of 8 for NCP and 9 for SUP out of 69 roosts in the NCSR (12% or 13% respectively, Tables 5, 6). Pelican numbers typically peak in the NCSR in fall, which is not included in the seasonal special closures, thus benefits to this species are less than would appear. Still, the NCRSG MPA Proposal represents an increase in protection to roosting seabirds over existing conditions (Proposal 0). Because offshore roost rocks are relatively abundant in the NCSR, protection of roost sites is far less critical than protection of large nesting colonies.

Nearshore Seabird Foraging Areas

The nearshore foraging analysis focused on four species with limited foraging ranges during the breeding season: Brandt's Cormorant, Common Murre, Pelagic Cormorant and Pigeon Guillemot. Weighted areas were calculated by multiplying seabird colony size as a percent of the bioregion population with the amount of that colony's foraging area captured by a given MPA. It is important to understand that this captures the amount of foraging area around colonies, so that special closures contribute little to this metric as they provide protection only to a small area around the breeding colonies themselves. Also, some of the state marine conservation areas (SMCAs) with certain allowable uses are not counted in this analysis because those uses diminish their contribution to these species (see Table 9.2 in *Draft Methods Used to Evaluate Marine Protected Area Proposals in the MLPA North Coast Study Region*). Table 7 shows the weighted area captured by each proposed MPA and special closure. Table 8 compares the Round 3 proposal and special closure recommendation based on the total weighted areas captured by MPAs and special closures that met the criteria for this analysis.

The NCRSG MPA Proposal increases benefits to nearshore foraging seabirds over Proposal 0. Because MPAs were generally not placed near major colonies (and special closures protect so little forage area), benefits to seabird foraging were tiny compared to the potential if MPAs were co-located close to colonies. Pyramid Point SMCA (not included in NCP, included in SUP), South Cape Mendocino SMR, and Ten Mile SMR were all close enough to colonies to achieve appreciable benefits (Table 7). Vizcaino SMCA also captured forage area but was not

included in NCP or SUP due to the proposed allowed uses. Pelagic Cormorants and Pigeon Guillemots are essentially obligate near shore foragers, thus improved foraging benefits to these species are more important than for Brandt's Cormorants and Common Murres which can range farther offshore.

Neritic Foraging Hot Spots

The neritic foraging analysis identified areas of persistent use by pelagic foraging seabirds and marine mammals and quantified the amount of these areas captured by proposed MPAs and special closures. Species groups were selected by their differing habitat use, thus 'hot spot' (defined as the top 10% of density in the NCSR) locations are likely to differ between groups. Table 9 shows the neritic hot spot areas captured by proposed MPAs and special closures from Round 3 and number of birds in 4 species groups using the area (but not including MPAs that did not overlap with a hot spot for any species group). Table 10 compares the total protected hot spot areas within SMRs, SMCAs that met the criteria for this analysis and special closures.

Considering the entire MPA array, Pyramid Point and Reading Rock SMCAs stood out as capturing hotspot areas for 3 species groups, and the Vizcaino SMCA and Ten Mile SMR captured hotspot area for the remaining group (mostly Common Murre and Brandt's Cormorant, Table 9). When only SMRs and qualifying SMCAs are considered, the Pyramid Point SMCA (in SUP) and Ten Mile SMR (in both NCP and SUP) were the only MPAs to protect hotspots (Table 10).

As with the near colony foraging analysis, it is important to understand that this analysis measures important foraging area at sea, and because special closures encompass little ocean surface, they contribute little to this analysis.

Shorebirds, Waterfowl, Estuarine Waterways and Coastal Habitats

The estuary and coastal habitats analysis quantified the amount of estuary, tidal flat, coastal marsh and beach habitat protected by proposed MPAs. All proposed special closures are located around offshore rocks and do not include any of these habitats, and are, therefore, not included in this analysis. Table 11 compares the species groups protected in estuaries in Round 3 MPAs. Data used for this analysis does not include estuaries south of the Eel River. The NCRSG MPA Proposal did not include SMRs or SMCAs that met the criteria to benefit these species groups, therefore there is no summary table of benefits. The NCRSG MPA Proposal also did not provide potential protection of shorebirds in Humboldt Bay because only a single state marine recreational management area (SMRMA), which allows waterfowl hunting, was proposed in Humboldt Bay. Therefore, no proposed MPAs met the criteria to benefit these species groups and no summary table of benefits was created.

The NCRSG MPA Proposal includes only a portion of south Humboldt Bay, that does include excellent foraging habitat for shorebirds and waterfowl, but since it also allows hunting, there will likely be a level of disturbance that reduces its value during the hunting seasons.

The Ten Mile Estuary SMRMA will provide some benefits to small numbers of estuarine waterfowl, mainly diving ducks and herons when waterfowl hunting is not occurring.

Summary

Seabird Breeding Colonies and Hot Spots

The NCRSG Special Closures Recommendation is very beneficial to seabirds, particularly at Castle Rock, one of the largest seabird colonies in the continental United States. Castle Rock is used by a great variety of birds and marine mammals throughout the year. The inclusion of Green and Flatiron Rocks north of Trinidad as seasonal special closures would have improved the special closures recommendation. The other breeding colony hot spot special closures are appropriately designated as seasonal. For nesting seabirds, the important segment of the year is between 1 March and 31 August. The NCRSG MPA Proposal and NCRSG MPA Proposal – Supplemental Evaluation without associated special closures includes very few important seabird breeding areas.

Seabird Roosting Sites

Unlike areas in other parts of the state, seabird roosting sites are common here on the north coast, and only a few of them are large, consistent roosts. Protection of important roost sites in the proposed MPAs and special closures represents an improvement over existing conditions.

Nearshore Foraging Areas

The benefits provided by protecting nearshore foraging areas are not as significant as the protection of breeding sites, but can benefit seabirds nonetheless. Because few MPAs in NCP and SUP were located close to breeding colonies, benefits to nearshore foraging species were not substantial, but still represent an improvement over existing conditions. This is particularly true for Pigeon Guillemots and Pelagic Cormorants, who depend on prey resources close to their nesting areas.

Neritic Foraging Areas

The 4 species groups were designated based on differing foraging habitat patterns, and consequently their foraging hot spots vary, making summary comparisons difficult. Pyramid Point SMCA (included only in SUP), Reading Rock SMCA (not included in either NCP or SUP), Rockport Rocks Seasonal Special Closures, and Vizcaino Rock Seasonal Special Closure overlapped with foraging hotspots. Larger protected areas would have provided more benefits to seabirds for neritic foraging.

Shorebirds, Waterfowl, Estuarine Waterways and Coastal Habitats

The NCRSG MPA Proposal will have minor, but positive benefits to shorebirds and waterfowl in southern Humboldt Bay (outside of the waterfowl hunting season) and slight positive benefits to herons and some waterfowl in the Ten Mile River estuary.

Special Closures

The SMCAs and SMRs currently proposed would provide protection only against consumptive activities. Non-consumptive activities such as close approach by boats, kayaking and surfing can still create disturbances at seabird breeding and roosting sites. Tremendous benefits to breeding seabirds can be provided using the special closures. Seasonal closures can provide excellent protection to breeding seabirds and so are recommended at the hot spots. Year round closures achieve the same results with the added protection to roosting birds and pinnipeds during the non-breeding season, such as the proposed special closure at Castle Rock.

Literature Cited

- Ainley, D. G., and R. J. Boekelheide (Eds.). 1990. Seabirds of the Farallon Islands: ecology, structure, and dynamics in an upwelling-system community. Stanford Univ. Press, Stanford, California.
- Ainley, D.G. W.J. Sydeman, and J. Norton. 1995. Upper-trophic level predators indicate interannual negative and positive anomalies in the California Current food web. *Mar. Ecol. Progr. Ser.* 118: 69-79.
- Anderson, D.W. and F. Gress. 1984. Pelicans and the anchovy fishery off southern California. *In* D.N. Nettleship, G.A. Sanger, and P.F. Springer eds. *Marine birds: their feeding ecology and commercial fisheries relationships*. Dartmouth, Nova Scotia: Canadian Wildlife Service Special Publication.
- Beale, C. M. and P. Monaghan. 2004. Human disturbance: people as predation-free predators? *Journal of Applied Ecology* 41:335-343.
- Bouton, S. N., P. C. Frederick, C. D. Rocha, A. T. Barbosa, and T. C. Bouton. 2005. Effects of tourist disturbance on Wood Stork nesting success and breeding behavior in the Brazilian Pantanal. *Waterbirds* 28:487-497.
- Cairns, D. K. 1992. Bridging the gap between ornithology and fisheries science: use of seabird data in stock assessment models. *Condor* 94:811-824.
- Carney, K.M., and W. J. Sydeman. 1999. A review of human disturbance effects on nesting colonial waterbirds. *Waterbirds* 22: 68-79.
- Clapp, R.B., M.K. Klimkiewicz, and J.H. Kennard. 1982. Longevity records of North American birds: Gaviidae through Alcidae. *Journal of Field Ornithology* 53(2): 81-124.
- Ford, R.G., D.G. Ainley, J.L. Casey, C.A. Keiper, L.B. Spear, and L.T. Ballance. 2004. The biogeographic patterns of seabirds in the central portion of the California Current. *Marine Ornithology* 32: 77-96.
- Golet, G.H., K.J. Kuletz, D.D. Roby, and D.B. Irons. 2000. Adult prey choice affects chick growth and reproductive success in Pigeon Guillemots. *Auk* 117(1): 82-91.
- Harris, J., M. Haward, J. Jabour, and E.J. Woehler. 2007. A new approach to selecting Marine Protected Areas (MPAs) in the Southern Ocean. *Antarctic Science* 19(2): 189-194.
- Jaques, D. and C. Strong. 2002. Disturbance to Brown Pelicans at communal roosts in southern and central California. Unpubl. report, Crescent Coastal Research, Astoria, Oregon.
- Jaques, D., C. S. Strong, and T. W. Keeney. 1996. Brown Pelican roosting patterns and responses to disturbance at Mugu Lagoon and other nonbreeding sites in the Southern California Bight. Technical Report No. 54, National Biological Service, Cooperative National Park Resources Studies Unit, Tuscon, Arizona.

- Massey, B.W., D.W. Bradley, and J.L. Atwood. 1992. Demography of a California Least Tern colony including effects of the 1982-1983 El Niño. *Condor* 94: 976-983.
- McChesney, G. J. 1997. Breeding biology of the Brandt's Cormorant on San Nicolas Island, California. Unpublished M.S. thesis, California State University, Sacramento, California.
- Monaghan, P. 1996. Relevance of the behavior of seabirds to the conservation of marine environments. *Oikos* 77: 227-237.
- Nur, N. and W. J. Sydeman. 1999. Survival, breeding probability and reproductive success in relation to population dynamics of Brandt's Cormorants *Phalacrocorax penicillatus*. *Bird Study* 46:S92-S103.
- Palmer, R.S. 1962. Brown Pelican. *In Handbook of North American Birds*, Vol. I. Yale University Press. New Haven. 567pp.
- Peters, K. A., and D. L. Otis. 2006. Wading bird response to recreational boat traffic: does flushing translate into avoidance. *Wildlife Society Bulletin* 34:1383-1391.
- Pichegru, L., P.G. Ryan, C. Le Bohec, C.D. van der Lingen, R. Navarro, S. Petersen, S. Lewis, J. van der Westhuizen, and D. Grémillet. 2009. Overlap between vulnerable top predators and fisheries in the Benguela upwelling system: implications for marine protected areas. *Marine Ecology Progress Series* 391: 199-208.
- Pierotti, R. and C.A. Annett. 1990. Diet and reproductive output in seabirds: food choices by individual, free-living animals can affect survival of offspring. *BioScience* 40(8): 568-574.
- Robinette, D.P., J. Howar, W.J. Sydeman, and N. Nur. 2007. Spatial patterns of recruitment in a demersal fish as revealed by seabird diet. *Marine Ecology Progress Series* 352: 259-268.
- Rojek, N. A., M. W. Parker, H. R. Carter and G. J. McChesney. 2007. Aircraft and vessel disturbance to Common Murres at breeding colonies in central California, 1997-1999. *Marine Ornithology* 35:67-75.
- Thayer, J.A., W. J. Sydeman, N. P. Fairman, and S. G. Allen. 1999. Attendance and effects of disturbance on coastal Common Murre colonies on Point Reyes, California. *Waterbirds* 22: 130-139.
- Safina, C. and J. Burger. 1988. Ecological dynamics among prey fish, bluefish, and foraging Common Terns in an Atlantic coastal system. *In* J. Burger, ed. *Seabirds and other marine vertebrates: competition, predation, and other interactions*. Columbia University Press. New York.
- Sydeman, W. J., M. M. Hester, J. A. Thayer, F. Gress, P. Martin and J. Buffa 2001. Climate change, reproductive performance and diet composition of marine birds in the southern California Current system, 1969-1997. *Progress in Oceanography* 49: 309-329.
- Wallace, E. A. H. and G. E. Wallace 1998. Brandt's Cormorant (*Phalacrocorax penicillatus*). *In* The Birds of North America, No. 362 (A. Poole and F. Gill, Eds.). The Birds of North America, Inc., Philadelphia, Pennsylvania.
- Warzybok, P. M., and R.W. Bradley. 2007. Population size and reproductive performance of seabirds on Southeast Farallon Island, 2007. Unpublished report, PRBO Conservation Science, Petaluma, California.
- Yen, P.P.W., W.J. Sydeman, and K.D. Hyrenbach. 2004. Marine bird and cetacean associations with bathymetric habitats and shallow-water topographies: implications for trophic transfer and conservation. *Journal of Marine Systems* 50: 79-99.

TABLES

Table 1. Numbers of breeding seabirds of 12 species within the north coast study region

Species	No. Animals
Total Number of Species	12
Black Oystercatcher (BLOY)	248
Brandt's Cormorant (BRCO) ^a	13105
Cassin's Auklet (CAAU)	4833
Common Murre (COMU)	258010
Double-crested Cormorant (DCCO)	2873
Fork-tailed Storm-Petrel (FTSP)	419
Leach's Storm-Petrel (LESP) ^b	9414
Pelagic Cormorant (PECO)	5675
Pigeon Guillemot (PIGU)	3148
Rhinoceros Auklet (RHAU)	1063
Tufted Puffin (TUPU)	181
Western Gull (WEGU)	4046
Study Region Total	303014

^a *American Ornithologists' Union (AOU) code for Brandt's Cormorant has been updated to BRAC since this data was collected.*

^b *American Ornithologists' Union (AOU) code for Leach's Storm-petrel has been updated to LHSP since this data was collected.*

Table 2. Numbers and percentages of marine birds at breeding colonies in Round 3 MPAs and special closures. ^a Not included in Table 3 because benefits to seabirds are reduced by allowed take activities.

Name	No. of Species	Total Birds (No.)	Total Birds (%)	BLOY	BRCO	COMU	DCCO	FTSP	LESP	PECO	PIGU	RHAU	TUPU	WEGU
Proposal 0														
(None in Proposal 0)														
NCP														
Pyramid Point SMCA ^a	4	52	<0.1%	3 (1.2%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	31 (0.5%)	12 (0.4%)	0 (0.0%)	0 (0.0%)	6 (0.1%)
South Cape Mendocino SMR	4	9690	3.2%	0 (0.0%)	464 (3.5%)	9163 (3.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (0.0%)	0 (0.0%)	0 (0.0%)	62 (1.5%)
Sea Lion Gulch SMR	2	19	<0.1%	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	17 (0.3%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	2 (0.0%)
Vizcaino SMCA ^a	4	46	<0.1%	3 (1.2%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	9 (0.2%)	20 (0.6%)	0 (0.0%)	0 (0.0%)	14 (0.3%)
Ten Mile SMR	5	525	0.2%	3 (1.2%)	257 (2.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	169 (3.0%)	58 (1.8%)	0 (0.0%)	0 (0.0%)	38 (0.9%)
SUP														
Pyramid Point SMCA	4	52	<0.1%	3 (1.2%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	31 (0.5%)	12 (0.4%)	0 (0.0%)	0 (0.0%)	6 (0.1%)
South Cape Mendocino SMR	4	9690	3.2%	0 (0.0%)	464 (3.5%)	9163 (3.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (0.0%)	0 (0.0%)	0 (0.0%)	62 (1.5%)
Sea Lion Gulch SMR	2	19	<0.1%	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	17 (0.3%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	2 (0.0%)
Vizcaino SMCA ^a	4	46	<0.1%	3 (1.2%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	9 (0.2%)	20 (0.6%)	0 (0.0%)	0 (0.0%)	14 (0.3%)

Name	No. of Species	Total Birds (No.)	Total Birds (%)	BLOY	BRCO	COMU	DCCO	FTSP	LESP	PECO	PIGU	RHAU	TUPU	WEGU
Ten Mile SMR	5	525	0.2%	3 (1.2%)	257 (2.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	169 (3.0%)	58 (1.8%)	0 (0.0%)	0 (0.0%)	38 (0.9%)
Special Closures														
Southwest Seal Rock Special Closure	4	151	0.0%	5 (2.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	134 (2.4%)	6 (0.2%)	0 (0.0%)	0 (0.0%)	6 (0.1%)
Castle Rock Special Closure	11	119796	39.5%	4 (1.6%)	2490 (19.0%)	108318 (42.0%)	0 (0.0%)	100 (23.9%)	926 (9.8%)	392 (6.9%)	360 (11.4%)	1005 (94.5%)	82 (45.3%)	1370 (33.9%)
False Klamath Rock Seasonal Special Closure	8	44980	14.8%	2 (0.8%)	713 (5.4%)	43898 (17.0%)	84 (2.9%)	0 (0.0%)	0 (0.0%)	115 (2.0%)	72 (2.3%)	0 (0.0%)	4 (2.2%)	92 (2.3%)
Sugarloaf Island Special Closure	8	1648	0.5%	3 (1.2%)	293 (2.2%)	0 (0.0%)	274 (9.5%)	0 (0.0%)	0 (0.0%)	627 (11.0%)	172 (5.5%)	7 (0.7%)	4 (2.2%)	268 (6.6%)
Steamboat Rock Seasonal Special Closure	4	9690	3.2%	0 (0.0%)	464 (3.5%)	9163 (3.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (0.0%)	0 (0.0%)	0 (0.0%)	62 (1.5%)
Rockport Rocks Seasonal Special Closure	7	2509	0.8%	1 (0.4%)	847 (6.5%)	1544 (6.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	91 (1.6%)	8 (0.3%)	2 (0.2%)	0 (0.0%)	16 (0.4%)
Vizcaino Rock Seasonal Special Closure	7	8799	2.9%	2 (0.8%)	1698 (13.0%)	6930 (2.7%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	70 (1.2%)	42 (1.3%)	1 (0.1%)	0 (0.0%)	56 (1.4%)

Note: Proposed MPAs and special closures not included in the table do not contain breeding seabird colonies.

Table 3. Comparison of numbers and percentages of marine birds breeding within Round 3 SMRs, qualifying SMCAs and special closures

Name	Black Oyster-catcher	Brandt's Cormorant	Common Murre	Double-crested Cormorant	Fork-tailed Storm-petrel	Leach's Storm-petrel	Pelagic Cormorant	Pigeon Guillemot	Rhinoceros Aukle	Tufted Puffin	Western Gull
P0	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
NCP	3 (1.2%)	721 (5.5%)	9163 (3.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	186 (3.3%)	59 (1.9%)	0 (0.0%)	0 (0.0%)	102 (2.5%)
SUP	6 (2.4%)	721 (5.5%)	9163 (3.6%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	217 (3.8%)	71 (2.3%)	0 (0.0%)	0 (0.0%)	108 (2.7%)
Special Closures	17 (6.9%)	6505 (49.6%)	169853 (65.8%)	358 (12.5%)	100 (23.9%)	926 (9.8%)	1429 (25.2%)	661 (21.0%)	1015 (95.5%)	90 (49.7%)	1870 (46.2%)

Table 4. Comparison of protection of the top eight marine bird breeding hot spots

Breeding Hot Spots	Proposal 0	NCP	SUP	Special Closures
Castle Rock				Castle Rock Special Closure
False Klamath Rock				False Klamath Rock Seasonal Special Closure
Green Rock				
Flatiron Rock				
False Cape Rocks				
Steamboat Rock		South Cape Mendocino SMR	South Cape Mendocino SMR	Steamboat Rock Seasonal Special Closure

Breeding Hot Spots	Proposal 0	NCP	SUP	Special Closures
Rockport Rocks				Rockport Rocks Seasonal Special Closure
Cape Vizcaino				Vizcaino Rock Seasonal Special Closure

Table 5. Brown Pelican roosts by roost size category within Round 3 MPAs and special closures. Not included in Table 6 because benefits to seabirds are reduced by allowed take activities.

MPA Name	Roost Category	Number of Roosts
Proposal 0		
MacKerricher SMCA ^a	Low	1
NCP		
Pyramid Point SMCA ^a	Low	1
Reading Rock SMCA ^a	Low	1
South Cape Mendocino SMR	Low	1
Vizcaino SMCA ^a	Low	2
Ten Mile SMR	Low	2
SUP		
Pyramid Point SMCA	Low	1
Reading Rock SMCA ^a	Low	1
South Cape Mendocino SMR	Low	1
Vizcaino SMCA ^a	Low	2
Ten Mile SMR	Low	2
Special Closures		
Castle Rock Special Closure	Low	1
False Klamath Rock Seasonal Special Closure	Medium	1
Steamboat Rock Seasonal Special Closure	Low	1
Rockport Rocks Seasonal Special Closures	Low	1
Vizcaino Rock Seasonal Special Closure	Low	1

Note: Proposed MPAs and special closures not included in the table do not contain Brown Pelican roosts.

Table 6. Comparison of size and number of Brown Pelican roosts within Round 3 SMRs, qualifying SMCAs and special closures

Draft MPA Proposal	High (>500 birds)	Medium (100-500 birds)	Low (never more than 100 birds)
Proposal 0	0	0	0
NCP	0	0	3
SUP	0	0	4
Special Closures	0	1	4

Table 7. Total contributions of nearshore weighted foraging index for four species of breeding seabirds in Round 3 MPAs and special closures. Not included in Table 8 because benefits to seabirds are reduced by allowed take activities.

Name	BRCO	PECO	COMU	PIGU	Name	BRCO	PECO	COMU	PIGU
Proposal 0					Special Closures				
Mackerricher SMCA ^a	0.00	<.01	0.00	0.02	Southwest Seal Rock Special Closure	0.00	<.01	0.00	<.01
Point Cabrillo SMCA ^a	0.02	0.02	<.01	0.02	Castle Rock Special Closure	0.01	<.01	0.03	<.01
Punta Gorda SMR	0.00	<.01	0.00	0.00	False Klamath Rock Seasonal Special Closure	<.01	<.01	<.01	<.01
Russian Gulch SMCA ^a	<.01	<.01	<.01	<.01	Sugarloaf Island Special Closure	<.01	<.01	<.01	<.01
Van Damme SMCA ^a	<.01	<.01	<.01	<.01	Steamboat Rock Seasonal Special Closure	<.01	<.01	<.01	<.01
					Rockport Rocks Seasonal Special Closure	<.01	<.01	<.01	<.01
					Vizcaino Rock Seasonal Special Closure	<.01	<.01	<.01	<.01
NCP					SUP				
Pyramid Point SMCA ^a	0.59	0.76	0.00	1.17	Pyramid Point SMCA	0.59	0.76	0.00	1.17
Point St. George Reef Offshore SMCA ^a	0.00	0.04	0.00	<.01	Point St. George Reef Offshore SMCA ^a	0.00	0.04	0.00	<.01
Reading Rock SMR	0.13	<.01	0.07	<.01	Reading Rock SMR	0.13	<.01	0.07	<.01
Reading Rock SMCA ^a	0.09	<.01	0.05	<.01	Reading Rock SMCA ^a	0.09	<.01	0.05	<.01
Samoa SMCA ^a	0.00	0.00	0.00	0.00	Samoa SMCA ^a	0.00	0.00	0.00	0.00
South Humboldt Bay SMRMA ^a	0.00	0.00	0.00	0.00	South Humboldt Bay SMRMA ^a	0.00	0.00	0.00	0.00
South Cape Mendocino SMR	0.28	0.53	0.17	0.26	South Cape Mendocino SMR	0.28	0.53	0.17	0.26
Mattole Canyon SMR	0.02	0.02	0.00	0.00	Mattole Canyon SMR	0.02	0.02	0.00	0.00
Sea Lion Gulch SMR	0.00	0.02	0.00	0.00	Sea Lion Gulch SMR	0.00	0.02	0.00	0.00
Vizcaino SMCA ^a	2.82	0.51	0.48	0.52	Vizcaino SMCA ^a	2.82	0.51	0.48	0.52
Ten Mile SMR	0.19	0.55	0.00	0.50	Ten Mile SMR	0.19	0.55	0.00	0.50
Ten Mile Beach SMCA ^a	0.05	0.08	0.00	0.05	Ten Mile Beach SMCA ^a	0.05	0.08	0.00	0.05
Ten Mile Estuary SMRMA ^a	<.01	<.01	0.00	<.01	Ten Mile Estuary SMRMA ^a	<.01	<.01	0.00	<.01
Point Cabrillo SMR	0.04	0.03	<.01	0.03	Point Cabrillo SMR	0.04	0.03	<.01	0.03

Name	BRCO	PECO	COMU	PIGU	Name	BRCO	PECO	COMU	PIGU
Big River Estuary SMP ^a	<.01	<.01	<.01	0.01	Big River Estuary SMP ^a	<.01	<.01	<.01	0.01
Navarro River Estuary SMRMA ^a	<.01	<.01	0.00	<.01	Navarro River Estuary SMRMA ^a	<.01	<.01	0.00	<.01

Note: MPAs and special closures not shown did not contribute to nearshore foraging area for any of these species.

Table 8. Comparison of draft MPA proposals to total contributions of weighted foraging areas for four species of breeding seabirds

	Brandt's Cormorant	Pelagic Cormorant	Common Murre	Pigeon Guillemot
Proposal 0	0.00	0.00	0.00	0.00
NCP	0.66	1.15	0.24	0.80
SUP	1.24	1.91	0.24	1.97
Special closures	0.03	0.02	0.04	0.02

Table 9. Comparison of diversity, area protected and mean number of birds contained in neritic foraging hot spots that overlap with Round 3 MPAs and special closures. ^a Not included in Table 10 because benefits to seabirds are reduced by allowed take activities.

MPA name	Species Diversity	Area (sq. mi.)	Average Number of Animals Sighted			
			Loons, Grebes and Scoters	Pigeon Guillemots and Pelagic Cormorants	Marbled Murrelets	All Other Seabirds
Proposal 0						
MacKerricher SMCA	13	0.5	-	9.15	-	-
NCP						
Pyramid Point SMCA ^a	17	8.27	287.5	88.5	197.7	-
Reading Rock SMCA ^a	14	7.73	2387.8	-	348.2	-
Vizcaino SMCA ^a	15	20.68	-	163.3	-	1961.4
Ten Mile SMR	14	8.57	-	-	-	1129.0
SUP						
Pyramid Point SMCA	17	8.27	287.5	88.5	197.7	-
Reading Rock SMCA ^a	14	7.73	2387.8	-	348.2	-
Vizcaino SMCA ^a	15	20.68	-	163.3	-	1961.4
Ten Mile SMR	14	8.57	-	-	-	1129.0
Special Closures						
False Klamath Rock Seasonal Special Closure	16	0.07	-	0.88	-	-

MPA name	Species Diversity	Area (sq. mi.)	Loons, Grebes and Scoters	Pigeon Guillemots and Pelagic Cormorants	Marbled Murrelets	All Other Seabirds
			Average Number of Animals Sighted			
Rockport Rocks: Seasonal Special Closure	8	0.01	-	-	-	2.36
Vizcaino Rock Seasonal Special Closure	8	0.01	-	-	-	1.54

Note: MPAs and special closures not shown did not contribute to neritic foraging hot spot area for any of these species.

Table 10. Comparison of total neritic foraging hot spot area protections for 4 species groups of seabirds within SMR and SMCA that meet protection criteria for seabirds

Name	Species Diversity	Area (sq. mi)	Loons, Grebes and Scoters	Pigeon Guillemots and Pelagic Cormorants	Marbled Murrelets	All Other Seabirds
			Average Number of Animals Sighted			
Proposal 0	-	-	-	-	-	-
NCP	14	13.85	-	-	-	1129.02
SUP	17	22.12	287.47	88.51	197.67	1129.02
Special Closures	16	0.23	-	0.88	-	3.91

Table 11. Comparison of protection of estuarine species groups and associated area of estuary in proposed MPAs.

MPA Name	% Area of Estuary in Proposed MPA	# of groups represented	Dabbling Ducks	Diving Ducks	Geese	Sea ducks	Shore-birds	Swans
			Proposal 0 (none in Proposal 0)					
NCP								
South Humboldt Bay SMRMA	7.18%	6	medium	high	high	high	high	high
SUP								
South Humboldt Bay SMRMA	7.18%	6	medium	high	high	high	high	high

Notes: MPAs not shown did not contribute to estuarine species protection. Data did not include estuaries south of the Eel River.

All state marine recreational management areas (SMRMAs) allow waterfowl hunting.

ATTACHMENT B

California MLPA Master Plan Science Advisory Team Evaluation of Benefits to Marine Mammals from Round 3 Proposed Marine Protected Areas and Special Closures *October 14, 2010*

The objective of this evaluation is to assess what benefits associated with goals 1, 2 and 4 of the California Marine Life Protection Act (MLPA) are achieved by proposed marine protected areas (MPAs) and special closures as they apply to marine mammals in the MLPA North Coast Study Region (NCSR). Proposed MPAs are evaluated for benefits for pinnipeds (seals and sea lions) and cetaceans (whales and porpoises). These animals are long-lived, produce few offspring and would benefit from placement of MPAs because of the reduction of disturbance from human activities.

Pinnipeds feed at sea and congregate onshore at traditional locations to rest at 'haulout sites' and to breed at 'rookeries'. These terrestrial sites are within intertidal or supratidal zones of the mainland and on islands. A range of substrates are represented at these sites, including hard rock, cobble and sand. Pinnipeds would benefit from the reduction of disturbance on or adjacent to rookeries or haulout sites. Vessel traffic, including motorized and non-motorized traffic, can cause significant levels of disturbance to marine mammals (e.g. Allen et al. 1985, Suryan and Harvey 1999, Thompson et al. 2001, Johnson and Acevedo-Gutierrez 2007). Disturbances can lead to reductions in productivity or site abandonment. Disturbances at foraging areas can disrupt feeding activities and cause animals to leave the area, further reducing feeding and leading to additional energy expenditures. Although MPAs do not restrict human access or vessel transit, the restrictions on allowable activities within MPAs are likely to result in fewer extractive users that access these areas. The proposed MPAs would provide protection only against consumptive activities. Non-consumptive activities such as kayaking and surfing can still create disturbances to marine mammals. This can be addressed through the use of no-entry special closure areas. Special closures are considered to provide the greatest benefit to marine mammals, followed by state marine reserves (SMRs).

Five pinniped species occur in the NCSR: Steller and California sea lions, northern fur seals, harbor seals, and elephant seals. Northern fur seals are rare and are generally seen offshore, and elephant seals only occur (and breed) at one location (Castle Rock) and are otherwise rare and found offshore. Species most likely to benefit include the two locally common and breeding pinnipeds, Steller sea lions and harbor seals, and the seasonally common and non-breeding California sea lion.

Most cetaceans (whales and dolphins) travel large distances and are not typically associated with a site that might be considered within the MPA framework. In the NCSR, summering gray whales and resident harbor porpoises are exceptions to the typical cetacean pattern as they are locally common and depend upon specific areas of the near-shore waters in this region. While gray whales typically migrate through the NCSR during the winter and spring, there is a small population that feed along our coast during the summer months. Northern California is the southern terminus of a subgroup of the gray whale population called the Pacific Coast Feeding Aggregation (PCFA) that forgo a full summer migration to Arctic seas and forage on benthic, epibenthic and swarming invertebrates along the coasts of Southeast Alaska, British Columbia, Washington, Oregon and northern California (NOAA, 2002). Harbor porpoise are locally resident and abundant in the nearshore waters throughout the year and are included in our analyses. Harbor porpoise breed and feed in nearshore waters. Gray whales and harbor

porpoise would benefit from the placement of MPAs because of the reduction of disturbance from human activities.

Methods

Evaluations follow the methods described in the *Draft Methods Used to Evaluate Marine Protected Area Proposals in the MLPA North Coast Study Region*. The evaluation includes analyses of the potential benefits to pinnipeds at: 1) breeding, 2) resting areas, and 3) nearshore foraging, and to 4) pinnipeds and cetaceans at neritic foraging areas.

Our analyses consider pinniped haulout sites, rookeries and forage areas that have been proposed with very high levels of protection (state marine reserves (SMR) or special closure areas) and do not include MPAs with lower levels of protection. We assume that most activities that affect pinnipeds on land would be reduced by these levels of protection. We recognize that protection of an area as a SMR does not address all potential sources of human activities, and that no-entry special closures would provide the highest level of benefit to marine mammals. We also recognize that lower levels of protection may provide some measure of protection. Data that directly evaluate potential impacts to pinnipeds in the NCSR are limited. Therefore these analyses provide a summary of the potential added value to pinnipeds due to proposed SMRs and special closures.

Population in this evaluation refers to the number of animals that use a site for breeding or resting. A haulout site is a location where seals and sea lions come onshore to rest. A rookery is a location where seals and sea lions come onshore to give birth, raise their young and breed. Many sites serve as both haulouts and rookeries.

Rookery and Hot Spot Analyses

For rookeries, or breeding sites, the two species most likely to benefit from MPAs include Steller sea lions and harbor seals. These species are sensitive to disturbance from human activities - particularly when breeding.

Analyses of pinniped rookery and haulout counts are drawn from survey data provided by Mark Lowry from NOAA Fisheries (pers. com.). Because harbor seal census data were collected just after pupping, during the molt period, systematic documentation of rookery locations in the NCSR is not available. We have conservatively characterized harbor seal haulouts of over 20 animals as rookeries.

There are two Steller sea lion rookeries and 62 harbor seal rookeries in the NCSR. Proposed MPAs and special closures that captured these rookeries were identified (Table 1). The two Steller sea lion rookeries were identified as hot spots due to their significance to the region and to the threatened eastern stock of Steller sea lions. Four hot spots were identified for harbor seal haulouts based on the large number of harbor seals that breed in these areas. Proposed MPAs and special closures that captured these hot spots were identified (Table 3).

Resting (Haulout) Sites

California sea lions, Steller sea lions and harbor seals will likely benefit from MPAs that protect haulout sites. Evaluation of the Round 3 MLPA North Coast Regional Stakeholder Group (NCRSG) MPA Proposal considers the total number of each species of pinniped, and calculates this as a percentage of the total number of pinnipeds within the NCSR for each proposed MPA and proposed special closure (Table 4). A comparison between Proposal 0 (existing MPAs), NCRSG MPA Proposal and the Round 3 NCRSG Special Closure Recommendation, with respect to the number of pinniped species, the number of each species of pinniped, and the percentage of the study region is presented in Table 5.

Foraging Areas

Harbor seals are the species most likely to benefit from potential increases to their forage base provided by MPAs. In nearshore areas, harbor seals typically forage near their haulout or rookery sites, and may repeatedly visit specific foraging areas (Jones 1981, Harvey and Torok 1994, Harvey et al. 1995, Thompson et al. 1998). Harbor seals forage on prey that is locally abundant, and they feed over a variety of habitats where they pursue rockfish, anchovies, squid and other prey (Table 9.1 in *Draft Methods Used to Evaluate Marine Protected Area Proposals in the MLPA North Coast Study Region*).

Steller sea lions are also likely to benefit during the breeding season from increases to their nearshore forage base provided by MPAs. During this time, adult females forage close to the rookery and consistently return to the rookery to care for their pups (Reimer et al., 2001).

To evaluate the effectiveness of MPAs to capture these foraging areas, buffers were created along three miles of coast and out to three miles offshore from haulouts and rookeries for harbor seals (rookeries only for Steller sea lions). These buffers were overlaid with proposed MPAs and special closures and the area of overlap determined. The proportion of the harbor seal and Steller sea lion foraging range overlapping proposed MPAs and special closures was then weighted based on the proportion of the study region population present within that proposed MPA or special closure. These weighted foraging indexes for SMRs and special closures for harbor seals (Table 6) and Steller sea lions (Table 7) are provided. The values are unitless but are useful to compare between proposals.

Neritic Foraging Areas

In addition to feeding near rookeries and haulouts, pinnipeds, whales and porpoises feed in other parts of nearshore waters. Neritic hot spots have been identified as places where pinnipeds, harbor porpoises and gray whales congregated during at sea systematic transect surveys (Strong, C., unpublished data). The neritic hot spots foraging analysis included plotting densities of these species over proposed MPAs and special closures to determine the area of neritic foraging hot spots protected for pinnipeds, harbor porpoises and gray whales. Hot spots were identified as areas with the top 10% of the density of observed pinnipeds (all species), harbor porpoises and gray whales (Appendix A). Our evaluation included the area of foraging 'hot spots' captured in proposed SMRs and special closures and the expected number of animals per area (Table 9).

In addition to the neritic transect surveys, gray whale foraging areas were also evaluated by plotting maximum densities of gray whales obtained from 12 years of shore-based surveys (Goley, P.D., unpublished data) over proposed SMRs and special closures. Five shore sites were identified from which whales were counted during the summer months: Point St. George, Crescent City Overlook, Klamath River mouth, Wedding Rock and Trinidad Bay and a three mile buffer was described around these sites. The proportion of the foraging range overlapping proposed MPAs and special closures was then weighted based on the average maximum count of whales during the summer months at each observation site in each MPA. The percentage of each observation area was then multiplied by the maximum number of whales counted at each site during the summer months (Table 8). The values are unitless but are useful for comparison between proposals.

Results

Three species of pinnipeds occur regularly in the north coast study region (California sea lion, Steller sea lion, harbor seal). Steller sea lions are locally abundant and are known to breed in the study region. There are two Steller sea lion rookeries in the study region (Sugarloaf Island and Southwest Seal Rock). These are biologically significant as they are two of the most southern rookeries of this threatened species. Harbor seals are also known to pup in the area and we estimate that there are 62 rookeries in the NCSR. Harbor seals are also locally abundant and known to breed in the region. California sea lions do not breed in the area, but are seasonally abundant on nearshore rocky haulouts.

Rookeries and Hot spots

Steller sea lions

The NCRSG MPA Proposal does not include any Steller sea lion rookeries within SMRs.

The NCRSG Special Closures Recommendation includes special closures that would protect the Steller sea lion rookeries on both Southwest Seal Rock and Sugarloaf Island year round (Tables 1 and 2). These two sites were also identified as pinniped hot spots (Table 3).

Harbor seals

The NCRSG MPA Proposal includes a very limited number (6 of 62) of harbor seal rookeries in proposed SMRs and special closures (Tables 1 and 2).

Of the 4 harbor seal rookeries that were identified as pinniped hot spots due to large numbers of breeding seals, one (vicinity of Castle Rock) was captured by the NCRSG MPA Proposal and the NCRSG Special Closures Recommendation. The mouth of the Eel River, Arcata Bay and South Humboldt Bay were not included in a proposed SMR or proposed special closure in Round 3. The NCRSG Special Closure Recommendation includes Castle Rock harbor seal rookery as a special closure, which does provide some protection for harbor seals.

Resting Sites

Analysis of Round 3 proposed SMRs and special closures

The NCRSG MPA Proposal includes four SMRs that provide benefits to a small portion of the NCSR population of harbor seals (1.5%), California sea lions (0.2%) and Steller sea lions (7.1%) at resting sites, which is less than many of the draft proposals in Round 2. The NCRSG Special Closures Recommendation also provides benefits to a small portion of the NCSR population of harbor seals (6.6%) and California sea lions (10.9%), and a large portion of the NCSR population of Steller sea lions (50.8%).

Nearshore Foraging Analysis

Harbor seals

The potential benefit from SMRs and special closures protecting likely foraging areas for Pacific harbor seals is summarized by the weighted foraging index (Table 6). Harbor seal rookeries and haulouts were included in only three SMRs in the NCRSG MPA Proposal. The 3 mile buffer around rookeries and haulouts intersected, to a very small extent, five of the proposed SMRs. This resulted in very low foraging indexes in the NCSR. Of the 1563 square miles identified as foraging areas, only 1.6% was captured by SMRs and special closures.

While proposed special closures benefit a small proportion of harbor seals within the area they surround with a 300 ft no entry zone, they provide a very limited benefit to the majority of foraging areas for harbor seals.

Steller sea lions

The potential benefit from proposed SMRs and special closures protecting likely foraging areas for Steller sea lions is summarized by the weighted foraging index (Table 7). Steller sea lion foraging indices were calculated for proposed SMRs. Given that no SMRs captured rookeries, there is little overlap between adjacent proposed SMRs leading to low foraging indices for Steller sea lions. Of the 47.4 square miles identified as foraging areas, only 10.1% was captured by SMRs and special closures.

While proposed special closures benefit Steller sea lions on the rock they surround with a 300 ft. no entry zone, they provide a very limited benefit to foraging areas for Steller sea lions.

Gray whales

The potential benefit from recommended special closures protecting foraging areas for gray whales during the summer months was negligible (weighted foraging index = 0.01). None of the buffered feeding areas were intersected by proposed SMRs in the NCRSG MPA Proposal (Table 8). Of the 99 square miles identified as foraging areas, only 0.07% was captured by SMRs and special closures.

Neritic foraging analysis

The potential benefits from MPAs and special closures protecting important foraging areas for all pinnipeds, harbor porpoise and gray whales based on at-sea transects was very low (Table 9 and Appendix A). When considering SMRs and special closures only, area of overlap with marine mammal neritic foraging areas was 5.51 square miles (about half what some of the Round 2 proposal achieved). The NCRSG MPA Proposal captured harbor porpoise foraging hot spots and only the False Klamath Rock Seasonal Special Closure intersected with gray whale hot spots. This analysis measures important foraging area at sea, and because special closures encompass little ocean surface, they contribute little to this analysis.

Summary

The NCRSG MPA Proposal and NCRSG Special Closures Recommendation include the addition of SMRs not currently in the network and special closures. Some of the SMRs and special closures proposed will benefit marine mammals by reducing disturbance at pinniped haulouts and rookeries. Additionally, the proposed networks provide very limited potential foraging benefits to marine mammals. The protection of Southwest Seal Rock and Sugarloaf Island Steller sea lion rookeries are noteworthy. These proposed special closures will provide tremendous benefit to the threatened Steller sea lions in the NCSR. When considering the combined impacts of the proposed SMRs and the special closures to the percentage of pinniped populations, the NCRSG MPA Proposal and the NCRSG Special Closures Recommendation offer some protection.

It is noteworthy that the NCRSG MPA Proposal does not provide a significant benefit to harbor seals with only 1.5% of the NCSR harbor seal population included in the NCRSG MPA Proposal and 6.6% included in the NCRSG Special Closures recommendation. Three of the harbor seal breeding hot spots were not included in NCRSG MPA Proposal and NCRSG Special Closures Recommendation. Harbor seals are considered one of the species most likely to benefit from MPAs and they play a significant role in the marine ecology of northern California. They reside in the NCSR year round, forage in the nearshore waters, and are dependent upon the local coastline and nearshore rocks to breed, molt and rest.

Pinniped and gray whale foraging areas did not significantly benefit from protection in the NCRSG MPA Proposal and NCRSG Special Closures Recommendation. We identified marine mammal foraging areas by creating 3 mile buffers around Steller sea lion and harbor seal rookeries as well as by identifying at-sea foraging hot spots for pinnipeds and cetaceans and identifying buffered zones around nearshore gray whale foraging areas. When considering proposed SMRs, the weighted foraging indexes were very low suggesting that the NCRSG MPA Proposal and NCRSG Special Closures Recommendation do not provide benefits to the foraging habitat that marine mammals in the NCSR depend upon. Steller sea lions, harbor seals, harbor porpoise and gray whales were identified as species most likely to benefit from MPAs, yet their foraging areas were largely missed by proposed MPAs.

Given that there are few SMRs that directly overlap with the identified marine mammal breeding or foraging hot spots, there is little protection to neritic or near shore marine mammal foraging hot spots offered in the NCRSG MPA Proposal and NCRSG Special Closures

Recommendation. In addition to the substrate required for breeding and resting, marine mammals are utterly dependent upon marine resources to feed. While special closures offer protection to pinnipeds on rookeries or haulouts, they contribute little to protection of the foraging areas. The 300 ft buffer currently proposed for special closures does offer some protection from direct disturbance to pinnipeds, which is very beneficial. However, this 300 ft buffer does not significantly contribute to protecting the foraging areas which are the 3 miles surrounding the rookeries/haulouts. These concerns would be addressed by including foraging areas within SMRs or modifying the special closures to include a no-entry zone of 1000 ft as has been implemented in other regions. These actions would contribute to increased weighted foraging indexes and would provide greater benefits to local marine mammals. While some rookeries and haulouts receive benefits from the NCRSG MPA Proposal, it has missed the opportunity to significantly improve benefits for harbor seals, specifically in protecting their foraging areas, or capture foraging areas for cetacean species over the existing MPAs.

Table 1. Comparison of number of rookeries within proposed MPAs by species and total number of rookeries within the north coast study region.

Name	Steller Sea Lion Rookeries	Harbor Seal Rookeries
Proposal 0		
MacKerricher SMCA ^a	0	3
Point Cabrillo SMCA ^a	0	1
NCRSG MPA Proposal		
Big Flat SMCA ^a	0	1
Vizcaino SMCA ^a	0	1
Ten Mile SMR	0	2
Point Cabrillo SMR	0	1
Special Closures		
Southwest Seal Rock Special Closure	1	0
Castle Rock Special Closure	0	1
Sugarloaf Island Special Closure	1	0
Rockport Rocks Seasonal Special Closure	0	1
Vizcaino Rock Seasonal Special Closure	0	1
Total Availability in Study Region	2	62

Note: Proposed MPAs and special closures not included in the table do not contain rookeries.

^a *Not included in Table 2 because benefits to marine mammals are reduced by allowed take activities.*

Table 2. Summary comparison of number of rookeries within proposed SMRs and special closures by species and total number of rookeries within the north coast study region.

Name	Total Pinniped Rookeries	Steller Sea Lion Rookeries	Harbor Seal Rookeries
Proposal 0	0	0	0
NCRSG MPA Proposal	3	0	3
Special Closures	5	2	3
Study Region Total	64	2	62

Table 3. Proposed special closures and SMRs containing pinniped population hot spots

Name	Population Hot Spots					
	SW Seal Rock	Sugarloaf Island	Vicinity of Castle Rock, Crescent City	South Bay, Humboldt Bay	Arcata Bay, Humboldt Bay	Mouth of the Eel River
Proposal 0						
NCP				South Humboldt Bay SMRMA ^a		
Special Closures	Southwest Seal Rock Special Closure	Sugarloaf Island Special Closure	Castle Rock Special Closure			

^a Designation does not provide benefits to marine mammals because of allowed take activities, but is included for reference only. Only special closure or SMR designation is considered to provide benefits to marine mammals.

Table 4. Number of animals and percentage of study region population within proposed MPAs and special closures

	Name	California Sea Lion	California Sea Lion %	Steller Sea Lion	Steller Sea Lion %	Harbor Seal	Harbor Seal %
Proposal 0	Point Cabrillo SMCA ^a	0	0.0%	0	0.0%	310	3.3%
	MacKerricher SMCA ^a	0	0.0%	0	0.0%	48	0.5%
NCP	Reading Rock SMCA ^a	3	<0.1%	4	0.1%	42	0.4%
	South Cape Mendocino SMR	25	0.2%	0	0.0%	14	0.2%
	Sea Lion Gulch SMR	0	0.0%	347	7.1%	0	0.0%
	Big Flat SMCA ^a	0	0.0%	16	0.3%	34	0.4%

	Name	California Sea Lion	California Sea Lion %	Steller Sea Lion	Steller Sea Lion %	Harbor Seal	Harbor Seal %
	Vizcaino SMCA ^a	54	0.4%	1	<0.1%	322	3.4%
	Ten Mile SMR	0	0.0%	0	0.0%	78	0.8%
	Point Cabrillo SMR	0	0.0%	0	0.0%	48	0.5%
Special Closures	Southwest Seal Rock Special Closure	5	<0.1%	1182	24.1%	0	0.0%
	Castle Rock Special Closure	1291	9.8%	716	14.6%	513	5.4%
	Sugarloaf Island Special Closure	56	0.4%	591	12.1%	20	0.2%
	Steamboat Rock Seasonal Special Closure	25	0.2%	0	0.0%	0	0.0%
	Rockport Rocks Seasonal Special Closure	0	0.0%	0	0.0%	66	0.7%
	Vizcaino Rock Seasonal Special Closure	58	0.4%	0	0.0%	25	0.3%

Notes: Proposed MPAs and special closures not included in the table do not contain pinniped haulouts or rookeries.

^a Designation does not provide benefits to marine mammals because of allowed take activities, but is included for reference only, and does not contribute to Table 5. Only special closure or SMR designation is considered to provide benefits to marine mammals.

Table 5. Comparison of number of species and number of animals at haul outs within proposed SMRs and special closures and total number of pinnipeds within the north coast study region.

	Number of Species	Animals (% of regional population)			Proposal Total Pinnipeds
		California Sea Lion	Steller Sea Lion	Harbor Seal	
Proposal 0	0	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
NCP	3	25 (0.2%)	347 (7.1%)	140 (1.5%)	512 (1.9%)
Special Closures	3	1435 (10.9%)	2489 (50.8%)	624 (6.6%)	4548 (16.5%)
Study Region Total	3	13200	4904	9451	27555

Table 6. Comparison of the harbor seal foraging index within proposed SMRs and special closures.

MPA Proposal	MPA Name	Weighted Forage Area	Sum of weighted area in SMRs
Proposal 0	Punta Gorda SMR	0.08	0.08
NCP	South Cape Mendocino SMR	0.05	1.17
	Mattole Canyon SMR	0.39	
	Sea Lion Gulch SMR	0.33	
	Ten Mile SMR	0.41	
	Point Cabrillo SMR	0.01	
Special Closures	Castle Rock Special Closure	<0.01	<0.01
	False Klamath Rock Seasonal Special Closure	<0.01	
	Sugarloaf Island Special Closure	<0.01	
	Steamboat Rock Seasonal Special Closure	<0.01	
	Rockport Rocks Seasonal Special Closure	<0.01	
	Vizcaino Rock Seasonal Special Closure	<0.01	

Table 7. Comparison of the Steller sea lion foraging index within proposed SMRs and special closures in the north coast study region.

	Name	Weighted Forage Area	Sum of weighted area in SMRs or special closures
Proposal 0	None	0.00	0.00
NCP	Point St. George Reef Offshore SMCA	0.44	0.58
	South Cape Mendocino SMR	0.58	
Special Closures	Southwest Seal Rock Special Closure	<0.01	0.01
	Steamboat Rock Seasonal Special Closure	<0.01	
	Sugarloaf Island Special Closure	<0.01	

Table 8. Gray whale foraging index within proposed SMRs and special closures

Name	SMR or Special Closure Name	Whales Weighted Forage Area
Proposal 0	None	0.00
NCP	None	0.00
Special Closures	Castle Rock Special Closure	0.01

Table 9. Neritic foraging hot spot area protection and number of animals at sea in proposed SMRs and special closures

Name	Area (sq. mi)	Average Number of Animals Sighted in Hot Spots		
		All Pinnipeds	Harbor Porpoise	Gray Whale
Proposal 0	-	-	-	-
NCP	5.28	55.1	-	-
Special Closures	0.23	2.2	-	0.1

Note: a dash indicates the area is not a hot spot for that species or group of species.

Literature Cited

- Allen, S. G., D. G. Ainley, G. W. Page, and C. A. Ribic. 1985. The effect of disturbance on harbor seal haul out patterns at Bolinas Lagoon, California, 1978-1979. U. S. Fishery Bull. 82: 493-500.
- Harvey, J. and M. Torok. 1994. Movements, dive behaviors, food habits of harbor seals (*Phoca vitulina richardsi*) in San Francisco Bay, California. Contract Number B-58227, California Department of Water Resources, Sacramento, CA.
- Harvey, J. R. Helm, G. Morejohn. 1995 Food habits of harbor seals inhabiting Elkhorn Slough, California. Calif Fish and Game 81:1-9.
- Johnson, A. and A. Acevedo-Gutierrez. 2007. Regulation compliance by vessels and disturbance of harbor seals. Canadian Journal of Zoology 85:290-294.
- Jones, R.E. 1981. Food habits of smaller marine mammals from northern California. Proc. Calif. Acad. of Sci., 42:409-433,
- Reimer, S. R. Brown, and B. Wright. 2001. The Steller Sea Lion in Oregon. International Marine Mammal conference, Vancouver.
- Suryan, R.M. and J.T. Harvey. 1999. Variability in reactions of Pacific harbor seals, *Phoca vitulina richardsi*, to disturbance. Fishery Bulletin 97:332-339.
- Thompson, P.M., S. Van Parijs and K.T. Kovacs. 2001. Local declines in the abundance of harbor seals: implications for the designation and monitoring of protected areas. Journal of Applied Ecology 38:117-125.
- Thompson, P.M., A. Mackay, D.J. Tollit, S. Enderby, and P.S. Hammond. 1998. The influence of body size and sex on the characteristics of harbor seal foraging trips. Canadian Journal of Zoology 76:1044-1053.

Appendix A

Table A1. Neritic foraging hot spot area and average number of animals at sea in proposed MPAs and special closures

Name	Area (sq. mi)	All Pinnipeds	Harbor Porpoise	Gray Whale
Average Number of Animals Sighted				
Proposal 0 (none)				
NCP				
Reading Rock SMCA	7.73	-	26.96	-
Samoa SMCA	15.67	-	82.16	-
South Cape Mendocino SMR	3.94	52.57	-	-
Mattole Canyon SMR	1.34	2.51	-	-
Vizcaino SMCA	20.68	30.74	-	-
Special Closures				
False Klamath Rock Seasonal Special Closure	0.07	-	-	0.09
Steamboat Rock Seasonal Special Closure	0.05	0.64	-	-
Sugarloaf Island Special Closure	0.09	1.24	-	-
Rockport Rocks Seasonal Special Closure	0.01	0.07	-	-
Vizcaino Rock Seasonal Special Closure	0.01	0.05	-	-

Note: Only MPAs with overlap of 'hot spot' foraging areas are included. A dash indicates the area is not a hot spot for that species or group of species.

^a *Designation does not provide benefits to marine mammals because of allowed take activities, and is not included in Table 9. Only SMRs and special closures, combined by proposal, are included in Table 9.*